A CANCER THERAPY RESULTS OF FIFTY CASES

SECOND EDITION

BY MAX GERSON, M.D.

TOTALITY BOOKS, PUBLISHERS

Del Mar, California

in many cases a valuable help in the therapy. A few patients cannot stand pancreatin; the majority are satisfied to have less digestive trouble with gas spasms and less difficulty in regaining weight and strength. We use the tablets after the detoxication; each contains five grains and is uncoated. The patient takes two or three tablets two or three times after meals, and later less.

It should always be borne in mind that cancer is a degenerative disease. The regeneration is only possible through the metabolism. Its restoration is hard work, but it is essential and

the last refuge for these advanced cases.

Retrospectively, I think the results were arrived at because I did not follow most of the scientific literature nor the laboratory findings, as far as they did not accord with the clinical confirmations. "Der Erfolg am Krankenbett ist entscheidend," Professor Kussmaul said. (The result at the sick-bed is decisive.) I do not want to make the mistake Winston Churchill expressed so clearly: "Men occasionally stumble over the Truth, but most pick themselves up and hurry off as if nothing had happened."

Rehabilitatio

THE TASK of rehabilitatic parable to that which he of cancer and the subset of rehabilitation is possible cases. Partial rehabilitaticases, and no rehabilitaticases, and no rehabilitaticases, and he goes home lowing problems:

(a) Medical

- I) The organization
- 2) Lack of help.
- Inability and iner type of diet.
- After weeks of pe iences and prepare out help or a doct
- Difficulty in shor foodstuffs.
- 6) Difficulties in pro drug stores (frequentes by pharmacis potassium glucona
- Re-examination—o

(b) Economic_

 Depletion of fund with previous treat

CHAPTER XXXI

Some Failures Medication: Not Specific

THE CREAT majority of my patients are far advanced cases who have little or no appetite and are barely able to take the necessary amount of juices and digest them. For that purpose gastric juice has to be applied in the form of acidol pepsin: two capsules three times daily before meals. In addition, they need the digestive enzymes of the pancreas as the poisoned organs stop or diminish the secretion: trypsin, lipase and diastase. All these are secreted by the pancreas; therefore, pancreatin is administered in some cases: three tablets three to five times a day, not during the first two weeks, and later less if needed. These enzymes are also needed for the so-called parenteral digestion of tumors and cancer: the natural activated enzymes are better, of course. The healing of cancer in the latter part of the treatment can be considered as a parenteral digestion. After I recognized the healing of cancer to be a parenteral digestion, the entire therapeutic endeavor was subordinated to this purpose. This means that after the cancer mass is killed, the dead piece must be dissolved; (I have a collection of such eliminated dead pieces from rectum, cervix, bladder, vagina, esophagus, tonsils, intestines, or wherever they find a way out of the body.)

The most drastic set-back occurred when I added to the therapy the opposite sex hormones, so highly recommended at that time by Professor Charles Higgins. These hormones were first applied in five cases which could not recover fast enough. In the first three to four months, I observed a pronounced improvement. Therefore, I administered these hormones to an additional 25 patients. All of these patients already were free of cancer



ecessary for strict coopera-

the dietary food as comig. nospitals or nursing homes, woid disturbance of family

uences.
some physicians.
toration of the entire body.
the present and the near

vill to live and to be cured. ten to twelve per cent of s of the disease; such realiw such a strict treatment. hen they see good results of pain in a matter of days. nent; some do not like to hers will not accept food ent problems. and pessimistic during the that they could not regain voman, who was with her : week; then the husband The patient was dismissed; but could not accomplish ividual cases can be found seases. ent and psychological co-

nment play important roles

patient needs faith, love,

slish this difficult task, the

self and favorable results

CHAPTER XXX

Most Frequent Mistakes of Patients in the Application of the Treatment

AFTER ABOUT four to six weeks in the clinic most of the patients feel stronger and have lost most of their original anxiety and depression. They think they can relax their adherence to rules and schedules. They frequently find no one to help them sufficiently with the preparation of the juices, diet, application of enemas, etc.

In general, people go to hospitals for operations or serious illness; the family considers them recovered upon their return. This is different with cancer. Cancer is a degenerative disease, not an acute one, and the treatment can be effective only if carried out strictly in accordance with the rules for one and a half to two years. We repeat here that it is not a symptom that is treated, nor a specific disease, but the reactions and functions of the entire body which have to be transformed and restored.)

As an illustration of the difficulties encountered, I quote the following from a patient's letter: "I feel like I have gone downhill since arriving home due to the strenuous past week—can't see how it could have been avoided. I have stayed on the diet—just have not managed to get in all the juices until the past two days." As an explanation of the foregoing, it should be mentioned that the patient had ten operations in eight years prior to arriving at the clinic. Naturally she was seriously weakened but gained strength in four weeks; this led her to the assumption that she could now take care of all the shopping, preparation and treatment by herself, without help.

Furthermore, at the time of their leaving home, patients often





trary I am of the opinion found effect on cancerous ecreased according to the composition."158

ing: the saltless diet and whole system. This is the with the reduction of the e way is paved for the minerals) and positively anges seem to force the n my opinion, the mineral number of other revived ole for the death of the only;) therefore they are hanges-they break down ust be properly composed ion of the liver. Thus, we ions, functions of the recation, etc., necessary for

ulating neoplastic growth cancer diet. 154 A contrary 1al and E. Hesse in 1935, or in salt have a rather ısias.

1-poor and K-rich regime or development, by induccancer without alkalosis." Berg,155 objected strongly calosis may be responsible hese opinions still remain

sodium and potassium in he authors and their conppointing. My opinion is 1as no uniform symptoms

d Diet, The Williams & Wilkins

and is not equally developed to a certain degree. Cancer is an extraordinary symptom only. The underlying cause is to be found in the poisoning of the liver. That is most probably the reason why the biological findings are inconclusive and so contradicting. Cancer is a disease of the liver lately called a "balance wheel of life"—where most metabolic functions are more or less concentrated. From here the other organs can be pathologically influenced and damaged or poisoned. Among the great number of observations, there are some which seem to be correct but they are not confirmed by laboratory experiments. Waterman found: "the sodium content of blood serum is unchanged in cancer patients." Benedict and Theis concluded that the "blood serum in cancer patients contains the normal amount of sodium." Pitts and Johnson examined the sodium content of blood serum and of blister fluid in cancerous and noncancerous patients and discovered that "the sodium content of these fluids was the same in cancer patients and in normal patients." Dr. Fry described in the British Cancer Review of 1926 the fact that in the blood of tumor-bearing rats the amount of sodium is 25 per cent above normal when the tumor is growing actively, and 60 per cent above normal when the tumor is receding.) Marwood went so far as to say salt is the root cause of cancer.

TASK OF THE SALTLESS DIET IN CANCER

The main task of the saltless diet is to eliminate the retained Na, Cl, H2O, together with toxins and poisons from the tissues all over the body.

All poisons and other substances difficult to eliminate are stimulants for the sick tissues, especially (liver and kidneys) That condition seems to be the reason why sodium chloride excretion increases in tuberculosis, cancer and other chronic diseases after two to three days on a saltless diet, and this condition stays at that higher level for about eight to ten or fourteen days, corresponding to a favorable development in the course of the disease. After that is accomplished, it stays near the normal level with the saltless diet, but shows a higher Na-Cl excretion, together with more fluid from time to time for two to three days, and later for one day. Such so-called "flare-ups" go along some-







times with nausea, diarrhea and nervous disturbances, caused probably by greater bile secretion and stimulation of the visceral nervous system. After each "flare-up" the patient feels easier and mentally improved.

Indications for Saltless Diet

(a) Edema and abnormal deposition of sodium and chloride in the subcutaneous tissue (nephropathias).

(b) Cardio-renal insufficiency.

(c) K-loss and Na-retention, in chronic diseases, especially in tuberculosis, cancer, etc.

(d) Detoxication, the degree of which must be in proportion to the degree of the disease—and which must be maintained during the period of restoration.

CHA

Ins

WE HAVE learned in recent insecticides is doing more a our bodies. I cannot emphasis represents our external metafield may read the Hearings. Investigate the Use of Chemaresentatives. Eighty-First C clearly described in the hear this field and what he recom

The following is a brief tion for uncontrolled genera DDT, or chlorophenothane substances that followed, h Beyond question, no other before developed so rapidl large a portion of the earth surprising as, at the time DI amount of data was alread showing that this agent w species of animals, that it fat and that it appeared in DDT poisoning in human observations were almost

"In the subsequent mas a vast amount of additions materials, both in animals

¹⁵⁶ Created Pursuant to H. J Nutritional Research, Milwaukee

y, consisting of two tablec coffee with brown sugar oil enema.

of enemas for regular use: caffeine drops from a ten of water of body temperanile extract and the prehe camomile extract, take ile flowers or leaves, or a of water. Let it boil for r ten minutes. Strain and vell covered, in the refrigied only in mild cases or

on take three tablespoons vater. Let it boil for three minutes or more. Strain he daily amount can be

itor oil treatment, the folike two large tablespoons c coffee, sweetened with ke a castor oil enema, as iter with toilet soap (no spoons of castor oil and i. Add 30 caffeine drops (consisting of de-fatted imals).

out rather a therapeutic the colon, we use half a e, as prepared for drinkflow in very slowly and s best when it is entirely re is a colostomy, we use iseased part, very slowly. ix, or urine bladder, we ese parts to help rid the body of odorous necrotic tissue discharges. Little bleedings are no contra-indication. This procedure is actually requested by patients as it brings them much relief from pain, discomfort and offensive odor.

Furthermore, it is necessary for the patient to drink freshly prepared vegetable juice every hour. This consists of four glasses of the juice of apples and carrots in equal parts, and also four glasses of green leaf juice. All these juices contain plenty of active oxidation enzymes enriched by a 10 per cent solution of minerals of the potassium group (potassium gluconate, potassium acetate, and potassium phosphate, monobasic). The oxidation enzymes of these juices, once pressed out of the cells and activated, are easily destroyed by oxygen from the air as well as from changes in light and temperature. They may lose 60 per cent of their active oxidation power within half an hour. Therefore, they must be consumed

immediately after pressing.

From the beginning, I felt that the tumor had to be killed while some scientists were satisfied to arrest the growths for as long a time as possible. In one of my articles I enumerated eleven points of difference between normal and cancer cells. The most important points are: cancer cells have (more Na) (ionised), live on (fermentation) (not on the normal oxidation), are negatively charged)electrically, do not have the normal exchange with blood and serum, and grow and spread uncontrolled. Studying these I felt there must be a way to prevent the fermentation, that is, to eliminate the basic facts upon which fermentation is built and can function. The fermentation is vital for the life of the cancer cell, That is the object upon which we could base further tests and explorations. How could it be done? The most Na-free diet has to be applied to extract Na from cancer cells through the blood and lymph stream. Instead of Na, potassium and the oxidizing enzymes have to be brought in with the help of an allergic inflammation. This reactivated power of the detoxified body had to be perfected to the highest degree as the cancer cells with their highly negative electrical potentials have the power to repulse forcefully whatever is counteracting their life process, maintained by fermentation.

The details have been explained elsewhere. We will concentrate on the parenteral digestion—the most important part for the

practice of the cancer treatment.



